

K-SOL



KEY POINTS

- Designed to provide a water-soluble source of vitamin K
- Contains menadione sodium bisulfite complex (MSBC), the best source of vitamin K for drinking water administration
- Pre-measured packet for error-free usage



DIRECTIONS:

Use 1 packet per 2 gallons* of stock solution for 5-7 days. Prior to management practices such as beak trimming or comb dubbing, K-Sol may be used at a rate up to 5 packets per 2 gallons of stock solution for 1 day.

** When used in proportioned set to deliver 1 oz per gallon of drinking water*

Poultry drinking water containing synthetic vitamin K should be kept out of direct sunlight.

Mix fresh solutions daily.

Store in a cool, dry place.

Ingredients: Sucrose, potassium chloride, sodium chloride, menadione sodium bisulfite complex.

Synthetic Vitamin K Soluble Crystals for use in Poultry Drinking Water

SKU: 3764

Zoetis and the Zoetis logo are registered trademarks of Zoetis, Inc.
Copyright © 2013 Zoetis, Inc.

5 Giralda Farms | Madison, NJ 07940 | (888) 963-8471 | zoetis.com

**FOR ORAL ANIMAL USE ONLY
NOT FOR HUMAN USE
KEEP OUT OF REACH OF CHILDREN**

**NET WEIGHT
188 g**

Contains: 10.6 g of MSBC per packet.

Take Time

**Observe Label
Directions**

zoetis™