

LINCOMYCIN SOLUBLE



KEY POINTS

- Potent antimicrobial for targeted activity against dysentery in swine or necrotic enteritis in broilers
- No withdrawal period
- Pre-measured packet for error-free usage
- Available in 2 convenient packet sizes



SWINE:

Lincomycin Soluble is indicated for the treatment of swine dysentery (bloody scours).

Dosage: Administer at a dose rate of 250 mg of lincomycin per gallon of drinking water. In clinical studies, this dose rate provided an average of 3.8 mg of lincomycin per lb of body weight per day.

Treatment Period: The drug should be administered for a minimum of 5 consecutive days beyond the disappearance of symptoms (bloody stools) up to a maximum of 10 consecutive days. If water treatment is discontinued prior to this time, a lincomycin treatment program may be continued with lincomycin premix at 100 g lincomycin per ton of complete feed as the sole ration according to label directions.

Administration: This packet will medicate 128 gallons of drinking water providing 250 mg/gallon. This packet will medicate 256 gallons of drinking water providing 250 mg/gallon for 160-gram packet. A dose of 3.8 mg lincomycin per lb of body weight may be maintained by medicating the drinking water at a concentration of 250 mg per gallon of drinking water when pigs are consuming 1.5 gallons per 100 lbs of body weight per day. Under these circumstances the concentration of lincomycin required in medicated water may be adjusted to compensate for variations in age and weight of animals, the nature and severity of disease symptoms, environmental temperature and humidity, each of which affects water consumption. For use in automatic water proportioners, prepare the stock solution by dissolving 1 packet in 1 gallon (2 gallons for 160 gram packet) of water: then adjust the proportioner to deliver 1 oz of stock solution per gallon of drinking water.

SKU: 3765 & 3766

Zoetis and the Zoetis logo are registered trademarks of Zoetis, Inc.
Copyright © 2013 Zoetis, Inc.

5 Giralda Farms | Madison, NJ 07940 | (888) 963-8471 | zoetis.com

BROILER CHICKENS:

Lincomycin Soluble is indicated for the control of necrotic enteritis caused by *Clostridium perfringens* susceptible to lincomycin.

Dosage: Administer at a dose rate of 64 mg of lincomycin per gallon of drinking water.

Treatment Period: Start medication as soon as the diagnosis of necrotic enteritis is determined. If improvement is not noted within 24 to 48 hours, consult a licensed veterinarian or veterinary diagnostic laboratory to determine diagnosis. The drug should be administered for 7 consecutive days.

Administration: This packet will medicate 500 gallons (1000 gallons for 150 gram packet) of drinking water providing 64 mg/gallon.

STORE BELOW 86° F (30° C).

Active Ingredient: Lincomycin HCl, equivalent to 32 g (64 g for 160 gram packet) lincomycin

FOR ORAL USE IN SWINE AND BROILER CHICKENS ONLY

KEEP OUT OF REACH OF CHILDREN

**NET WEIGHT 2.82 oz (80 g)
NET WEIGHT 5.64 oz (160 g)**

CAUTIONS:

1. Discard medicated drinking water if not used within 2 days. Fresh stock solution should be prepared daily.
2. If clinical signs of bloody scours (watery, mucoid or bloody stools) have not improved during the first 6 days of medication, discontinue treatment and redetermine the diagnosis.
3. Occasionally, swine fed lincomycin may within the first 2 days after the onset of treatment develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within 5 to 8 days without discontinuing the lincomycin treatment.
4. The safety of lincomycin has not been demonstrated for pregnant swine or swine intended for breeding.
5. Do not allow rabbits, hamsters, guinea pigs, horses or ruminants access to water containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.
6. Do not use the water treatment and the feed treatment simultaneously.
7. Not for use in layer and breeder chickens.

WARNINGS:

1. No drug withdrawal period is required before slaughter of swine receiving Lincomycin Soluble at the approved level of 250 mg per gallon of drinking water, nor before slaughter of birds receiving Lincomycin Soluble at the approved level of 64 mg per gallon of drinking water.
2. Not for human use.



zoetis[™]