

ALBON[®]

(sulfadimethoxine)

Boluses

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION: Albon is a low-dose, rapidly absorbed, long-acting sulfonamide.

Actions: Sulfadimethoxine has been demonstrated in laboratory studies to be effective against a wide variety of organisms, such as streptococci, staphylococci, and members of the *E. coli*-salmonella group of bacteria.

Comparatively low doses of Albon give rapid, sustained blood levels required for effective disease therapy.

INDICATIONS AND USAGE: Albon is effective in the treatment of shipping fever complex and bacterial pneumonia associated with *Pasteurella* spp. sensitive to sulfadimethoxine; and calf diphtheria and foot rot associated with *Fusobacterium necrophorum* sensitive to sulfadimethoxine in cattle.

WARNINGS:

Not for Human Use

Milk taken from animals during treatment and for 60 hours (5 milkings) after the latest treatment must not be used for food.

Do not slaughter animals for food purposes within 7 days following the last treatment.

A withdrawal period has not been established for this product in preruminating calves.

Do Not Use in Calves to be Processed for Veal



DISCARD MILK FOR

60 HOURS

AFTER LAST TREATMENT



DO NOT SLAUGHTER COW FOR

7 DAYS

AFTER LAST TREATMENT

PRECAUTIONS: During treatment period, make certain that animals maintain adequate water intake.

If animals show no improvement within 2–3 days, reevaluate your diagnosis. Treatment should not be continued beyond 5 days.

DOSAGE AND ADMINISTRATION: Albon should be administered to cattle so that the initial dose is equivalent to 25 mg/lb of body weight and each subsequent daily dose is equivalent to 12.5 mg/lb of body weight. Length of treatment will depend on clinical response. In most cases, treatment for 3–4 days is adequate. Treatment should not be continued beyond 5 days.

The following tables show the dosage and dosage forms to be used for cattle of different weights.

Albon Boluses, 5 g

Dosage Schedule for Cattle: 200–600 lb of body weight

Animal Weight, lb	First Day	Daily for the Following 3–4 Days
200	1 bolus	1/2 bolus
300	1 1/2 boluses	1 bolus
400	2 boluses	1 bolus
500	2 1/2 boluses	1 1/2 boluses
600	3 boluses	1 1/2 boluses

Albon Boluses, 15 g

Dosage Schedule for Cattle: 600–1,200 lb of body weight

Animal Weight, lb	First Day	Daily for the Following 3–4 Days
600	1 bolus	1/2 bolus
800	1 1/2 boluses	1 bolus
1,000–1,200	2 boluses	1 bolus

TOXICITY AND SAFETY: Albon has been shown to be a well-tolerated sulfonamide with relatively high solubility at the pH normally occurring in the kidney and with a low degree of toxicity. Following the administration of Albon at the recommended dosage, no undesirable side effects have been observed.

Store at Controlled Room Temperature 15°–30°C (59°–86°F)

HOW SUPPLIED: Albon Boluses are supplied in the following dosage forms:

5 g sulfadimethoxine per single-scored bolus

15 g sulfadimethoxine per single-scored bolus

Approved by FDA under NADA # 031-715

zoetis

Distributed by:

Zoetis Inc.,

Kalamazoo, MI 49007

40039314A&P
Revised: June 2022