

# Bovine Rhinotracheitis-Virus Diarrhea-Parainfluenza<sub>3</sub>- Respiratory Syncytial Virus Vaccine

Modified Live Virus

## Leptospira Canicola-Grippotyphosa- Hardjo-Icterohaemorrhagiae-Pomona Bacterin

### Bovi-Shield GOLD® FP® 5 L5



**INDICATIONS:** Bovi-Shield GOLD FP 5 L5 is for vaccination of healthy cows and heifers prior to breeding to prevent persistently infected calves caused by bovine virus diarrhea (BVD) virus Types 1 and 2 and as an aid in preventing abortion caused by infectious bovine rhinotracheitis (IBR, bovine herpesvirus Type 1) virus; respiratory disease caused by IBR, BVD Types 1 and 2, parainfluenza<sub>3</sub> (PI<sub>3</sub>) and bovine respiratory syncytial virus (BRSV); BVD Type 2 testicular infection; and leptospirosis caused by *Leptospira canicola*, *L. grippotyphosa*, *L. hardjo*, *L. icterohaemorrhagiae*, and *L. pomona*. A 12-month duration of immunity has been demonstrated against IBR-induced abortion and persistently infected calves caused by BVD Types 1 and 2. Bovi-Shield GOLD FP 5 L5 may be administered to pregnant cattle provided they were vaccinated, according to label directions, with any Bovi-Shield GOLD FP or PregGuard® GOLD FP vaccine within the past 12 months. Bovi-Shield GOLD FP 5 L5 may also be administered to calves nursing pregnant cows provided their dams were vaccinated within the past 12 months as described above. **To help ensure safety in pregnant cattle, heifers must receive at least 2 doses of any Bovi-Shield GOLD FP or PregGuard GOLD FP product with the second dose administered approximately 30 days prebreeding.**

**PRODUCT DESCRIPTION:** Bovi-Shield GOLD FP 5 L5 is a freeze-dried preparation of modified live virus (MLV) strains of IBR, BVD (Types 1 and 2), PI<sub>3</sub>, and BRSV viruses, plus a liquid bacterin containing the 5 *Leptospira* serovars identified above. The liquid bacterin is used to rehydrate the freeze-dried vaccine.

#### DIRECTIONS:

**General Directions:** Vaccination of healthy cattle is recommended. Aseptically rehydrate the freeze-dried vaccine (Bovi-Shield GOLD FP 5) with the liquid bacterin provided (Leptoform-5®+A), shake well, and administer 2 mL intramuscularly. In accordance with Beef Quality Assurance guidelines, this product should be administered in the muscular region of the neck.

**Primary Vaccination:** Administer a single 2-mL dose to all breeding cows and heifers approximately 1 month prior to breeding or being added to the herd.

**Revaccination:** Annual revaccination with a single dose of Bovi-Shield GOLD FP 5 L5 is recommended.

Good animal husbandry and herd health management practices should be employed.

#### PRECAUTIONS:

**Do not use in pregnant cows (abortions can result) unless they were vaccinated, according to label directions, with any Bovi-Shield GOLD FP or PregGuard GOLD FP vaccine within the past 12 months. Do not use in calves nursing pregnant cows unless their dams were vaccinated within the past 12 months as described above.**

**To help ensure safety in pregnant cattle, heifers must receive at least 2 doses of any Bovi-Shield GOLD FP or PregGuard GOLD FP product with the second dose administered approximately 30 days prebreeding.**

Store at 2°–7°C. Prolonged exposure to higher temperatures and/or direct sunlight may adversely affect potency. Do not freeze.

Use entire contents when first opened.

Sterilized syringes and needles should be used to administer this vaccine.

Do not sterilize with chemicals because traces of disinfectant may inactivate the vaccine.

Burn containers and all unused contents.

Do not vaccinate within 21 days before slaughter.

Contains gentamicin as preservative.

As with many vaccines, anaphylaxis may occur after use. Initial antidote of epinephrine is recommended and should be followed with appropriate supportive therapy.

This product has been shown to be efficacious in healthy animals. A protective immune response may not be elicited if animals are incubating an infectious disease, are malnourished or parasitized, are stressed due to shipment or environmental conditions, are otherwise immunocompromised, or the vaccine is not administered in accordance with label directions.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services, (888) 963-8471 (USA), (800) 461-0917 (Canada).

For veterinary use only.

U.S. Veterinary License No. 190

**zoetis**

Zoetis Inc.  
Kalamazoo, MI 49007, USA