

zoetis



12+
WEEKS



RABIES 1 YEAR

Injectable

50 Vials (1 dose: 1 mL)

Rabies Vaccine

KILLED VIRUS



- Annual revaccination
- For use in dogs, cats and ferrets only.

The logo for Vanguard, featuring a blue and red swoosh above the word "VANGUARD" in bold blue capital letters.



RABIES 1 YEAR Injectable

This product has been shown to be effective for the vaccination of healthy dogs, cats and ferrets 12 weeks of age or older against rabies. Duration of immunity of at least 1 year has been demonstrated after repeat dose. For more information regarding efficacy and safety data, go to productdata.aphis.usda.gov.

The vaccine is prepared from cell-culture-grown, chemically inactivated rabies virus. The seed virus is a highly immunogenic, fixed strain of rabies virus which originated from Louis Pasteur's original isolate in 1882. The inactivated virus is formulated with a highly purified adjuvant and is packaged in liquid form.

Directions: Shake well. Aseptically administer 1 mL subcutaneously. Dogs may be vaccinated intramuscularly or subcutaneously. A repeat dose should be administered 1 year later.

Precautions: Store at 2°–8°C. Prolonged exposure to higher temperatures may adversely affect potency. Do not freeze. Sterilized syringes and needles should be used to administer this vaccine. Contains gentamicin and thimerosal as preservatives. This product has not been tested in pregnant animals. As with many vaccines, anaphylaxis may occur after use. Initial antidote of epinephrine is recommended and should be followed with appropriate supportive therapy. Do not mix with other products. In case of human exposure, contact a physician.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services, (888) 963-8471.

This product has been shown to be efficacious in healthy animals. A protective immune response may not be elicited if animals are incubating an infectious disease, are malnourished or parasitized, are stressed due to shipment or environmental conditions, are otherwise immunocompromised, or the vaccine is not administered in accordance with label directions.