

RofenAid[®]

40

**Caution: Federal law restricts
medicated feed containing this
veterinary feed directive (VFD)
drug to use by or on the order
of a licensed veterinarian.**

**Sulfadimethoxine & Ormetoprim
Type A Medicated Article**

**Antibacterial
Anticoccidial**

Net wt 50 LB (22.68 kg)

zoetis

RofenAid[®] 40



RofenAid[®]

40

**Sulfadimethoxine & Ormetoprim
Type A Medicated Article**

**Antibacterial
Anticoccidial**

**For use in broiler, replacement (breeder and layer)
chickens, turkeys, ducks and chukar partridges only.**

Active drug ingredients	Sulfadimethoxine 113.5 grams per pound (25%), Ormetoprim 68.1 grams per pound (15%) in a carrier suitable for incorporation in feed.
For use in manufacturing medicated feeds only	
Broiler and replacement (breeder and layer) chickens	Indications: As an aid in the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. brunetti</i> and <i>E. mivati</i> ; as an aid in the prevention of bacterial infections caused by <i>Haemophilus gallinarum</i> (infectious coryza), <i>Escherichia coli</i> (colibacillosis) and <i>Pasteurella multocida</i> (fowl cholera). Use directions: Add 1 lb of RofenAid 40 to one ton of feed to provide 0.02% (113.5 g/ton [0.0125%] sulfadimethoxine and 68.1 g/ton [0.0075%] ormetoprim) concentration in feed. Feed continuously as the sole ration. Warning: Do not feed to chickens over 16 weeks (112 days) of age.
Turkeys	Indications: As an aid in the prevention of coccidiosis caused by <i>Eimeria adenoeides</i> , <i>E. gallopavonis</i> , and <i>E. meleagrimitis</i> ; for the prevention of bacterial infections caused by <i>Pasteurella multocida</i> (fowl cholera). Use directions: Add 1/2 lb of RofenAid 40 to one ton of feed to provide 0.01% (56.75 g/ton [0.00625%] sulfadimethoxine and 34.05 g/ton [0.00375%] ormetoprim) concentration in feed. Feed continuously as the sole ration.
Ducks, including breeding ducks	Indications: As an aid in the control of bacterial infections due to <i>Pasteurella multocida</i> (fowl cholera) in ducks including breeding ducks. Use directions: Add 2 lbs of RofenAid 40 to one ton of feed to provide 0.04% (227 g/ton [0.025%] sulfadimethoxine and 136.2 g/ton [0.015%] ormetoprim) concentration in feed. Medication should be started at the first signs of infection. Feed for 7 days as the sole ration.
Ducks	Indications: As an aid in the control of bacterial infections due to <i>Escherichia coli</i> , <i>Riemerella anatipestifer</i> (strains 2,3 & 4) and severe challenge of <i>P. multocida</i> (fowl cholera). Use directions: Add 4 lbs of RofenAid 40 to one ton of feed to provide 0.08% (454 g/ton [0.050%] sulfadimethoxine and 272.4 g/ton [0.030%] ormetoprim) concentration in feed. Medication should be started at the first signs of infection. Feed for 7 days as the sole ration. Caution: Not for breeding ducks. The safety of feeding sulfadimethoxine and ormetoprim at levels higher than 0.025% and 0.015%, respectively, in breeding ducks has not been established.
Chukar Partridges	Indications: For the prevention of coccidiosis caused by <i>Eimeria kofoidi</i> and <i>E. legionensis</i> . Use directions: Add 1 lb of RofenAid 40 to one ton of feed to provide 0.02% (113.5 g/ton [0.0125%] sulfadimethoxine and 68.1 g/ton [0.0075%] ormetoprim) concentration in feed. Feed continuously as the sole ration to young birds up to 8 weeks of age. Caution: The safety of feeding RofenAid 40 to breeding stock has not been established.
Warning	Do not feed to turkeys and ducks producing eggs for food. Withdraw 5 days prior to slaughter of broiler and replacement chickens, turkeys and ducks.
Note: Manufacture of Type B or C feeds from this product requires a Medicated Feed Mill License. Labels for feeds containing RofenAid 40 must contain appropriate indications, limitations and warnings as well as required feed ingredient information.	
Restricted Drug (California) - Use only as directed. Not for human use. Approved by FDA under NADA # 040-209 Store below 25°C (77°F), excursions permitted to 37°C (99°F).	
zoetis Distributed by: Zoetis Inc. Kalamazoo, MI 49007 Product of China	<div><div>FPO: UPC</div><div> 0 67219 12004 0</div></div> <div>Take Time  Observe Label Directions</div>
Net wt 50 LB (22.68 kg)	
10010602	