HOW TO IMPLANT WITH SYNOVEX C

(progesterone and estradiol benzoate implants)

100 mg progesterone and 10 mg estradiol benzoate per implant

PELLETS

Study the following instructions carefully, then proceed step by step, until the technique becomes routine. Many head can be implanted per hour by an experienced team, one member of which should be assigned to do nothing but the implantation. This person should maintain hand cleanliness and use sanitary instruments only.

-2-

HOW SUPPLIED

100 Dose

SYNOVEX® C Implants

Synovex C (progesterone and estradiol benzoate implants) is recommended for use in suckling beef calves up to 400 lbs of body weight. Synovex C is also recommended for improvement in rate of weight gain in steers weighing greater than 400 pounds and fed in confinement for slaughter when used as part of a re-implant program in which an initial Synovex C implant is followed at approximately 70 days by Synovex S. Do not use in yeal calves. Effectiveness and animal safety in yeal calves have not been established.

Do not use in bull calves intended for reproduction, or in calves less than 45 days old.

Store at controlled room temperature 20° - 25°C (68° - 77°F) with excursions between 15°-30°C (59°-86°F). Avoid excessive heat or humidity.

Not for use in humans. For animal use only Implant pellets in the ear only. Any other location is in violation of federal law. Do not attempt salvage of implanted site for human or animal food

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for yeal.

CAUTION


Bulling, rectal and vaginal prolapse, udder development, ventral edema and elevated tailheads may be reported in calves administered Synovex C implants

Keep this and all drugs out of the reach of children.

Restricted Drug (California)— Use Only as Directed.

SYNOVEX® EXACT 10® Applicator

SYNOVEX® SX10 Applicator



- 5 -

SYNOVEX® Metal Revolver



STEP 1 Loading the Applicator

following the directions outlined in the instruction manual accompanying each applicator.

Load the applicator

Additional information for the individual applicator user guides can be found at: www.growwithsynovex.com.

STEP 2 Restraint



The animal must be confined in a restraint mechanism (squeeze chute or head gate). The implant site on the back of the ear should be prepared by scrubbing with a firm bristled brush that has been soaked in a germicidal solution.

Note: If implanting horned cattle, greater safety is provided when the head is controlled by the use of a bull lead (nose tongs)

STEP 3 Implant Site

Divide the ear into three imaginary sections as illustrated. The implanted pellets should be deposited in the center one-third of the ear as shown. To accomplish this, the applicator needle should be inserted in the **outer**

one-third of the ear as indicated by the "X" in the illustration. Implanting too close to the head may cause abnormal sexual behavior. Care should be taken to avoid severing the major arteries of the


STEP 4 Insert Needle



Grasp the ear with one hand. While holding the applicator firmly with the other hand, penetrate the skin at the point shown by the "X". Thrust the needle under the skin taking care not to penetrate the car-


tilage. Ease the applicator forward (toward the base of the ear) until the full needle length is beneath the skin.

STEP 5 Pellet Implantation



When the needle is completely inserted, activate the instrument by squeezing the trigger completely. Refer to instructions for individual applicator.

STEP 6 Inspection



Check the implant site. If properly administered, the implants should lie in a straight line under the skin.

Re-cock the applicator. Disinfect the applicator needle in a germicidal

solution. The applicator is

now ready to implant the next animal.

Zoetis Inc. Kalamazoo, MI 49007

zoetis Distributed by:

100 mg progesterone and 10 mg estradiol benzoate per implant

CALF IMPLANTS

For Increased Rate of Weight Gain

This product was manufactured by a non-sterilizing process.

Do Not Use In Veal Calves Not For Use In Humans For Animal Use Only

zoetis

Approved by FDA under NADA # 009-576

40031049