zoetis

DAPi

6+ WEEKS

INJECT

DAPi

Injectable

25 Vials (1 dose)

STERILE DILUENT 25 Vials (1 mL)

Canine Distemper-Adenovirus Type 2-Parainfluenza Vaccine

MODIFIED LIVE VIRUS

Thimerosal Free

To Reconstitute:
Rehydrate the vaccine with the sterile diluent for vaccination.

VANGUARD®

VANGUARD®

DAPi Injectable

This product has been demonstrated to be effective for vaccination of healthy dogs 6 weeks or age or older against canine distemper virus (CDV), canine adenovirus type 1 (CAV1), canine adenovirus type 2 (CAV2), and canine parainfluenza virus (CPIV). Duration of immunity has not been established. For more information regarding efficacy and safety data, see productdata.aphis.usda.gov.

Vaccination and revaccination according to label directions have been demonstrated (under field conditions) to result in serum antibody titers that persist for 12-48 months against CDV (serum neutralization [SN] titer ≥ 1:32), CAV1 (SN ≥ 1:16), CAV2 (SN ≥ 1:16) and CPIV (SN ≥ 1:16).

Directions: Aseptically rehydrate the freeze-dried vaccine with the sterile diluent provided, shake well, and administer 1 mL subcutaneously or intramuscularly. Healthy dogs should receive 2 doses administered 3–4 weeks apart. The presence of maternal antibody is known to interfere with the development of active immunity in puppies and additional boosters will be required in most young animals. Historically, annual revaccination with this product was recommended. The need for annual booster vaccinations has not been established. For advice on revaccination, consult your veterinarian or the manufacturer.

Precautions: Store at 2°–8°C. Prolonged exposure to higher temperatures and/or direct sunlight may adversely affect potency. Do not freeze. Sterilized syringes and needles should be used to administer this vaccine. Do not sterilize with chemicals because traces of disinfectant may inactivate the vaccine. Inactivate unused contents before disposal. Contains gentamicin as preservative. This product has not been tested in pregnant animals. In case of anaphylactoid reaction, administer epinephrine. Do not mix with other products, except as specified above. In case of human exposure, contact a physician.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services, (888) 963-8471.

This product has been shown to be efficacious in healthy animals. A protective immune response may not be elicited if animals are incubating an infectious disease, are malnourished or parasitized, are stressed due to shipment or environmental conditions, are otherwise immunocompromised, or the vaccine is not administered in accordance with label directions.