

10 mL

10 Doses

Encephalomyelitis-West Nile Virus Vaccine

Eastern & Western, Killed Virus

For vaccination of healthy horses 10 months of age or older as an aid in the prevention of viremia caused by West Nile virus and as an aid in the prevention of equine encephalomyelitis due to Eastern and Western viruses.

West Nile-Innovator® + EW

For Veterinary Use Only

U.S. Vet. License No. 190

Zoetis Inc.

Kalamazoo, MI 49007, USA

zoetis

Store in the dark at 2° to 7°C (35° to 45°F). AVOID FREEZING. SHAKE WELL. Do not vaccinate within 21 days before slaughter.

Serial No.:

Exp. Date:

50161000

↓ Lift here for complete directions ↓

West Nile-Innovator® + EW

SAFETY: Field safety testing has demonstrated that this product is safe for use in horses 3 months of age or older. Efficacy has not been established in foals 3 months of age.

EFFICACY: Efficacy of the West Nile fraction was demonstrated in 10-month-old horses that received two doses of the vaccine and were challenged one year post-vaccination with West Nile virus. The vaccine protected 95% of the vaccinates against viremia, while 82% of the controls were viremic.

DOSE: Horses, inject one 1 mL dose intramuscularly using aseptic technique. Administer a second 1 mL dose 3 to 4 weeks after the first dose. A 1 mL booster dose should be given annually. Early revaccination may be advisable when horses are faced with an outbreak or with other conditions which might make heavy exposure likely.

CAUTION: Use entire contents when first opened. In some instances, transient local reactions may occur at the injection site. In case of anaphylactoid reaction, administer epinephrine.

Thimerosal, neomycin and polymyxin B added as preservatives.