

1 mL

1 Dose

## Encephalomyelitis-West Nile Virus Vaccine

Eastern & Western & Venezuelan, Killed Virus

## Tetanus Toxoid

### West Nile-Innovator<sup>®</sup> + VEWT

For intramuscular vaccination of healthy horses 10 months of age or older as an aid in the prevention of viremia caused by West Nile virus, and as an aid in the prevention of equine encephalomyelitis due to Eastern, Western and Venezuelan viruses, and tetanus.

Store in the dark at 2° to 7°C (35° to 45°F). AVOID FREEZING.  
SHAKE WELL. Do not vaccinate within 21 days before slaughter.

Thimerosal, neomycin and polymyxin B added as preservatives.

See reverse side for complete directions.

U.S. Vet. License No. 190

Zoetis Inc.  
Kalamazoo, MI 49007, USA

Serial No.:  
Nº de Lote:  
Expiration Date:  
Fecha de Venc.:

**zoetis**

50024600

West Nile-Innovator<sup>®</sup> + VEWT

↑ PEEL HERE  
↑ DESPEGUE AQUÍ

## **West Nile-Innovator® + VEWT**

**INDICATIONS:** For intramuscular vaccination of healthy horses 10 months of age or older as an aid in the prevention of viremia caused by West Nile virus, and as an aid in the prevention of equine encephalomyelitis due to Eastern, Western and Venezuelan viruses, and tetanus.

**EFFICACY:** Efficacy of the West Nile fraction was demonstrated in 10-month-old horses that received two doses of the vaccine and were challenged one year post-vaccination with West Nile virus. The vaccine protected 95% of the vaccinates against viremia, while 82% of the controls were viremic.

**SAFETY:** Field safety testing has demonstrated that this product is safe for use in horses 3 months of age or older. Efficacy has not been established in foals 3 months of age.

**DOSAGE AND ADMINISTRATION:** Horses, inject one 1 mL dose intramuscularly using aseptic technique. Administer a second 1 mL dose 3 to 4 weeks after the first dose. A 1 mL booster dose should be given annually. Early revaccination may be advisable when horses are faced with an outbreak or with other conditions which might make heavy exposure likely.

**CAUTION:** Store in the dark at 2° to 7°C (35° to 45°F). AVOID FREEZING. SHAKE WELL. Use entire contents when first opened. In some instances, transient local reactions may occur at the injection site. Do not vaccinate within 21 days before slaughter. In case of anaphylactoid reaction, administer epinephrine.

Thimerosal, neomycin and polymyxin B added as preservatives.

For Veterinary Use Only

**Remove serial number and expiration date label and place on the animal's medical chart.**