

10 mL

10 Doses

Encephalomyelitis-Rabies-West Nile Virus Vaccine

Eastern & Western, Killed Virus

Tetanus Toxoid

This product has been shown to be effective for the vaccination of healthy horses 10 months of age or older against rabies, West Nile virus, tetanus, and encephalomyelitis caused by Eastern and Western viruses. A duration of immunity of at least 14 months has been demonstrated against rabies and at least 12 months against

CORE EQ INNOVATOR™

VLN 190/PCN 48R5.21

Zoetis Inc.
Kalamazoo, MI 49007, USA

zoetis

West Nile virus. The duration of immunity against the remaining diseases is unknown. For more information regarding efficacy and safety data, go to productdata.aphis.usda.gov.

Store in the dark at 2°-8°C (35°-46°F). Do not freeze. Do not vaccinate within 21 days before slaughter.

Serial No.:

Exp. Date:

40021024

↓ Lift here for complete directions ↓

SAFETY: Field safety testing has demonstrated that this product is safe for use in horses 3 months of age or older. Efficacy of the West Nile, Eastern, and Western virus fractions and tetanus has not been established in foals 3 months of age.

DOSAGE AND ADMINISTRATION: Shake well. Aseptically administer a single 1-mL dose intramuscularly, followed by a single dose of West Nile-Innovator® + EWT 3-4 weeks later. Vaccinate against rabies 1 year later, and annually after that. For routine revaccination frequency against the other diseases, or advice on revaccination in the face of stress or exposure, consult your veterinarian or the manufacturer.

CAUTION: Use entire contents when first opened. This product has not been tested in pregnant animals. Contains gentamicin, neomycin, polymyxin B, thimerosal, and formalin as preservatives. In case of anaphylactoid reaction, administer epinephrine. Do not mix with other products. In case of human exposure, contact a physician.

For veterinary use only in accordance with state and local regulations.

Technical inquiries should be directed to Zoetis Inc. Technical Services, (888) 963-8471.