

Swine Influenza Vaccine, H1N1 & H1N2 & H3N2

Killed Virus

250-dose vial of vaccine

Rehydrate to 500 mL

500 mL vial of sterile diluent




zoetis




For use in swine only
For veterinary use only



U.S. Veterinary License No. 190
Zoetis Inc.
Kalamazoo, MI 49007, USA



zoetis



Ser
Exp

IMPRINT AREA
NO COPY
NO VARNISH

For vaccination of healthy swine, including pregnant sows and gilts, 3 weeks of age or older as an aid in preventing respiratory disease caused by swine influenza A viruses H1N1, H1N2 and H3N2. A duration of immunity of at least 10 weeks has been established against swine influenza A virus H3N2. This product contains the following Influenza A Virus isolates: A/Swine/Iowa/110600/2000 (H1N1); A/Swine/Oklahoma/0726H/2008 (H1N2); A/Swine/North Carolina/394/2012 (H3N2); and A/Swine/Minnesota/872/2012 (H3N2).

A sterile diluent containing Amphigen® is used to rehydrate the freeze-dried vaccine.

Directions: Shake diluent before use. Aseptically rehydrate the freeze-dried vaccine with the accompanying adjuvant-containing sterile diluent, shake well, and administer 2 mL intramuscularly. Healthy swine 3 weeks of age or older should receive two 2-mL doses administered approximately 3 weeks apart. In young pigs, vaccinate after maternally derived antibodies to swine influenza A viruses have declined. Semi-annual revaccination with a single dose is recommended.

Precautions: Store at 2°–7°C. Do not freeze. Use entire contents when first opened or rehydrated. Sterilized syringes and needles should be used to administer this vaccine. Do not vaccinate within 21 days before slaughter. Contains gentamicin as preservative. As with many vaccines, anaphylaxis may occur after use. Initial antidote of epinephrine is recommended and should be followed with appropriate supportive therapy.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services, (888) 963-8471 (USA), (800) 461-0917 (Canada).

Code 19A5.2C