

Porcine Circovirus Vaccine Type 1-Type 2 Chimera

Killed Virus

PCV2a and PCV2b

Contains
two PCV2
genotypes

This product has been shown to be effective for the vaccination of healthy pigs 3 days of age or older against porcine circovirus 2 (PCV2). A 23-week duration of immunity has been demonstrated. For more information regarding efficacy and safety data, go to productdata.aphis.usda.gov. This product contains PCV2a and PCV2b, and has been shown to be effective against PCV2 viremia, shedding of virus in feces, lymphoid colonization and lymphoid depletion.

Directions: Shake well. Aseptically administer a single 2-mL dose intramuscularly to healthy pigs 3 weeks of age or older. Fostera Gold PCV may also be given as two 1-mL doses administered three weeks apart to pigs 3 days of age or older. The optimal choice of the dosing regimen, including age at time of vaccination, will depend on farm circumstances.

Precautions: Store at 2°-8°C (35°-46°F). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. This product has not been tested in pregnant animals. Contains gentamicin and thimerosal as preservatives. The vaccine may induce mild to moderate transient local injection site reactions following vaccination in 3-day-old piglets. In case of allergic reaction, administer epinephrine. Do not mix with other products. In case of human exposure, contact a physician.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services,
(888) 963-8471.

For Veterinary Use Only
VLN 190/PCN 19K5.R1

Zoetis Inc., Kalamazoo, MI
49007, USA