



100 mL

Porcine Circovirus Vaccine, Type 1 – Type 2 Chimera

Killed Virus

Mycoplasma Hypopneumoniae Bacterin

PCV2a, PCV2b, and M. hyo

Contains
two PCV2
genotypes

This product has been shown to be effective for the vaccination of healthy pigs, including pregnant sows and gilts, 3 days of age or older against porcine circovirus 2 (PCV2) and respiratory disease due to *M. hyopneumoniae*. A

23-week duration of immunity has been demonstrated. For more information regarding efficacy and safety data, go to productdata.aphis.usda.gov. This product contains PCV2a and PCV2b, and has been shown to be effective against PCV2 viremia, shedding of virus in feces, lymphoid colonization and lymphoid depletion.

Directions: Shake well. Aseptically administer a single 2-mL dose intramuscularly to healthy pigs 3 weeks of age or older. Fostera Gold PCV MH may also be given as two 1-mL doses administered three weeks apart to pigs 3 days of age or older. The optimal choice of the dosing regimen, including age at time of vaccination, will depend on farm circumstances.

Precautions: Store at 2°-8°C. Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Contains gentamicin and thimerosal as preservatives. The vaccine may induce mild to moderate transient local injection site reactions following vaccination in 3-day-old piglets. In case of allergic reaction, administer epinephrine. Do not mix with other products. In case of human exposure, contact a physician.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services,

(888) 963-8471.

For Veterinary Use Only

VLN 190/PCN 49K5.R7

Zoetis Inc.

Kalamazoo, MI 49007, USA

The Zoetis logo, featuring the word "zoetis" in a lowercase, bold, sans-serif font. The letter "z" is stylized with a long, sweeping underline that extends to the right.