

For vaccination of healthy pigs 3 weeks of age or older as an aid in preventing viremia, lymphoid depletion, and colonization of lymphoid tissue caused by porcine circovirus type 2; and as an aid in reducing PCV2 virus shedding. A 23-week duration of immunity has been demonstrated.

Directions: Shake well. Aseptically administer a single 2-mL dose intramuscularly.

Fostera PCV with MetaStim may also be given as two 1-mL doses administered 2 weeks apart.

For Veterinary Use Only

500 mL

250 doses

Porcine Circovirus Vaccine Type 1-Type 2 Chimera

Killed Virus

zoetis

PCV
MetaStim®

Fostera™
PCV MetaStim®

Precautions: Store in the dark at 2°-7°C (35°-45°F). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Contains gentamicin and thimerosal as preservatives. Allergic reactions, although rare, may occur. The antidote is epinephrine.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services, (888) 963-8471 (USA), (800) 461-0917 (Canada).

U.S. Veterinary License No. 190

Zoetis Inc.
Kalamazoo, MI 49007, USA



50205300

Ser:
Exp:

Time/Date: 50205300.pdf 1 8/18/16 3:53 PM

zoetis		Project No.	Artwork DIR No.	Component Material No.	Description	Country			
		14328	50205300	40014388	Fostera PCV w/MetaStim	US			
		Dimensions		Drawing No.	FG Material No.	DIR Type/Description			
		8.5" x 3.5"		Z14-800210	10016568	Z64/Label			
Additional Info:		Colors:				<input type="checkbox"/> EDITOR'S COPY			
							GS: DATE:		
GS	T. Norman	Rev	GA	PR	CHANGES	GS / ART REV (LCA)	CHANGES	GS / ART REV (FA)	CHANGES
GA	L. Nelson	1			OK		OK		OK