

For vaccination of healthy pigs 3 weeks of age or older as an aid in preventing viremia, lymphoid depletion, and colonization of lymphoid tissue caused by porcine circovirus type 2; and as an aid in reducing PCV2 virus shedding and enzootic pneumonia caused by *M. hyopneumoniae*. A 23-week duration of immunity has been demonstrated.

**Directions:** Shake well. Aseptically administer a single 2-mL dose intramuscularly. Foster PCV MH may also be given as two 1-mL doses administered 2 weeks apart.

100 mL

50 doses

# Porcine Circovirus Vaccine, Type 1 – Type 2 Chimera

Killed Virus  
**Mycoplasma  
Hyopneumoniae Bacterin**

zoetis

PCV MH

**Precautions:** Store at 2°-8°C (35°-46°F). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. This product has not been tested in pregnant animals. Contains gentamicin and thimerosal as preservatives. Allergic reactions, although rare, may occur. The antidote is epinephrine. Do not mix with other products. In case of human exposure, contact a physician.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services, (888) 963-8471.

For Veterinary Use Only

VLN 190/PCN 49K5.R5

Zoetis Inc.  
Kalamazoo, MI 49007, USA

 **Foster**™  
PCV MH