

zoetis



Kalamazoo, MI 49007, USA

Zoetis Inc.

U.S. Vet. License No. 190

Thimerosal added as preservative.

Store in the dark at 2°-7°C. Avoid freezing. Shake well. Do not vaccinate within 21 days before slaughter.

serovar *pomona*.

For vaccination of healthy horses 6 months of age or older as an aid in the prevention of leptospirosis caused by *Leptospira interrogans*

Lepto EQ Innovator®

Leptospira Pomona Bacterin

1 dose

1 mL

1 mL

Leptospira Pomona Bacterin

1 dose

Lepto EQ Innovator®

For vaccination of healthy horses 6 months of age or older as an aid in the prevention of leptospirosis caused by *Leptospira interrogans* serovar *pomona*.

Store in the dark at 2°-7°C. Avoid freezing. Shake well. Do not vaccinate within 21 days before slaughter.

Thimerosal added as preservative.

U.S. Vet. License No. 190

Zoetis Inc.

Kalamazoo, MI 49007, USA

Serial No.

Exp. Date:

zoetis 50155000

Lepto EQ Innovator®

FEEL HERE

Lepto EQ Innovator®

INDICATIONS: For vaccination of healthy horses 6 months of age or older as an aid in the prevention of leptospirosis caused by *Leptospira interrogans* serovar *pomona*. Lepto EQ Innovator contains inactivated whole cultures of *L. pomona* with MetaStim adjuvant to enhance the immune response.

Efficacy: Immunogenicity was demonstrated by vaccination and challenge tests in 6-month-old, healthy horses. Thirty horses were vaccinated with 2 doses of either Lepto EQ Innovator (vaccinates) or saline (controls), followed by experimental challenge 3 weeks later with a heterologous strain of *L. pomona*. *L. pomona* was reisolated from all 15 control horses from at least 1 sample (urine, kidney and/or liver) while no *L. pomona* was reisolated from any sample from vaccinated horses. Ten weeks later, 100% of control animals were positive for leptospiremia, while no vaccinates were positive. Similarly, 93% of controls were positive for leptospiruria, while no vaccinates tested positive.

Safety: Field safety testing has demonstrated that this product is safe for use in horses 3 months of age or older, including pregnant mares. Efficacy has not been demonstrated in 3-month-old foals.

Dosage and administration: Horses, inject one 1-mL dose intramuscularly using aseptic technique.

Administer a second 1-mL dose 3-4 weeks after the first dose. Annual revaccination with a single dose is recommended.

Lepto EQ Innovator®

Caution: Shake well. Use entire contents when first opened. Contains thimerosal as preservative. In case of anaphylactoid reaction, administer epinephrine.

For Veterinary Use Only

Remove serial number and expiration date label and place on the animal's medical chart.