



# Porcine Epidemic Diarrhea Vaccine

## Killed Virus

This product license is conditional.  
Safety and efficacy studies are in progress.

50 doses    100 mL

zoetis

For vaccination of healthy pregnant sows and gilts pre-farrowing against diarrheal disease in their neonatal pigs caused by porcine epidemic diarrhea virus.

**Directions:** Shake well and administer 2 mL intramuscularly. Healthy pregnant swine should receive 2 doses with the first dose being administered 5 weeks prior to farrowing. The second dose should be given 2 weeks prior to farrowing. Pregnant swine should be vaccinated with a single dose 2 weeks before each subsequent farrowing. Duration of immunity has not been evaluated.

**Precautions:** Store at 2°-7°C. Do not freeze. Use entire contents when first opened. Sterilized syringes and needles should be used to administer this vaccine. Do not vaccinate within 21 days before slaughter. Transient site swelling may occur following vaccination. Contains gentamicin and thimerosal as preservatives. As with many vaccines, anaphylaxis may occur after use. Initial antidote of epinephrine is recommended.

For veterinary use only

Technical inquiries should be directed to Zoetis Inc. Veterinary Services, (888) 963-8471 (USA), (800) 461-0917 (Canada).

U.S. Veterinary License No. 190

Zoetis Inc., Kalamazoo, MI 49007, USA

30393401

Ser  
Exp