

**Easy to administer.**

**Indications and Usage:** For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. RFD Liquid Wormer may also be used to prevent reinfestation of *T. canis* in puppies and adult dogs and in lactating bitches after whelping.

**Shake Well Before Use**

**Active Ingredient:** 2.27 mg of pyrantel base as pyrantel pamoate per mL

**Recommended Storage:** Store below 30°C (86°F).

**Warning:** Keep out of reach of children.

**Dosage and Administration:** Administer 1 teaspoon (5 mL) for each 5 lb of body weight. To assure proper dosage, weigh animal prior to treatment. It is not necessary to withhold food prior to treatment.

Distributed by:  
Zoetis Inc.  
Kalamazoo, MI 49007



40028073  
Product of Spain

# RFD<sup>®</sup>

(pyrantel pamoate  
oral suspension)

## LIQUID WORMER

### FOR PUPPIES AND DOGS

Approved by FDA under NADA # 100-237

Net Contents:  
2 fl oz (60 mL)

Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2–4 weeks after first treatment. If your dog looks or acts sick, consult your veterinarian before treatment.

For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8 and 10 weeks of age. Lactating bitches should be treated 2–3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation.



Lot No.:

Exp. Date: