

Precautions: Store at 2°–7°C. Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. As with many vaccines, anaphylaxis may occur after use. Initial antidote of epinephrine is recommended and should be followed with appropriate supportive therapy.

Directions: Aseptically administer 2 1/2 mL subcutaneously, followed by a second dose 4–6 weeks later. Annual revaccination with a single dose is recommended.

Directions: Shake well. *Cattle:* Aseptically administer 5 mL subcutaneously, followed by a second dose 4–6 weeks later. In accordance with Beef Quality Assurance guidelines, this product should be administered subcutaneously (SC) under the skin. *Sheep:* Aseptically administer 2 1/2 mL subcutaneously, followed by a second dose 4–6 weeks later. Annual revaccination with a single dose is recommended.

Directions: Shake well. *Cattle:* Aseptically administer 5 mL subcutaneously, followed by a second dose 4–6 weeks later. In accordance with Beef Quality Assurance guidelines, this product should be administered subcutaneously (SC) under the skin. *Sheep:* Aseptically administer 2 1/2 mL subcutaneously, followed by a second dose 4–6 weeks later. Annual revaccination with a single dose is recommended.

Ultrabac® 7


For veterinary use only

Clostridium Chauvoei- Septicum- Novyi-Sordellii- Perfringens Types C & D Bacterin-Toxoid

50 cattle doses
100 sheep doses



250 mL



Technical inquiries should be directed to
Zoetis Inc. Technical Services,
(888) 963-8471 (USA), (800) 461-0917 (Canada).
U.S. Veterinary License No. 190

zoetis

Zoetis Inc.
Kalamazoo, MI 49007, USA

Ser
Exp



YES

250 mL
Ultrabac® 7

Administer 5 mL under the skin
on the neck.
Administrar 5 mL bajo la piel
en el cuello.

Remove Tab and
Place on Syringe
Barrel