

Canine Adenovirus Type 2- Parainfluenza- Bordetella Bronchiseptica Vaccine

MODIFIED LIVE VIRUS & AVIRULENT LIVE CULTURE

- Administer a single 0.5-mL dose intranasally.
- Store at 2°-8°C. Do not freeze.
- See back of card for complete directions and precautions.



zoetis

RAPID RESP 3

Intranasal (Syringe-Free)

25 Vials (1 dose)

STERILE DILUENT

25 Applicators (0.5 mL)

To Reconstitute:

Rehydrate the vaccine
with the sterile diluent
for vaccination.

**VANGUARD®**



RAPID RESP 3

Intranasal (Syringe-Free)

This product has been shown to be effective for the vaccination of healthy dogs 8 weeks of age or older against canine adenovirus type 2 (CAV2), canine parainfluenza virus (CPiV), and *B. bronchiseptica*. A duration of immunity of at least one year has been demonstrated against CAV2, CPiV, and *B. bronchiseptica*. For more information regarding efficacy and safety data, see productdata.aphis.usda.gov.

Directions: Remove tear-away seal and stopper from vial containing freeze-dried vaccine. Remove plastic tip from applicator, dispense sterile diluent into the vial and gently swirl. Draw contents back into the applicator and administer a single 0.5-mL dose intranasally. Historically, annual revaccination with this product was recommended. The need for annual booster vaccinations has not been established for this product; consultation with a veterinarian or the manufacturer is recommended.

Precautions: Store at 2°-8°C. Prolonged exposure to higher temperatures and/or direct sunlight may adversely affect potency. Do not freeze. Inactivate unused contents before disposal. This product has not been tested in pregnant animals. In case of anaphylactoid reaction, administer epinephrine. Do not mix with other products, except as specified above. In case of human exposure, contact a physician.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services, (888) 963-8471.