

FOALS



ANTHELCIDE[®]
EQ PASTE
(oxibendazole)

STRONGID[®]
PASTE
(pyrantel pamoate)



BABIES NEED SPECIAL DEWORMING CARE

A TARGETED APPROACH FOR CONTROLLING PARASITES DEPENDS ON WHEN YOUR FOAL WAS BORN

		AGE	2 MONTHS	4-6 MONTHS (Prior to weaning)	7-9 MONTHS (Approx. 1 month post-weaning)	9 MONTHS	12 MONTHS
EARLY FOALS BORN MARCH OR EARLIER 	TARGET PARASITES	ROUNDWORMS (ascarids)	ROUNDWORMS (ascarids)	STRONGYLES ROUNDWORMS* <i>(Only if detected via FEC)</i>	STRONGYLES & TAPEWORMS	STRONGYLES	
	DEWORMER	STRONGID [®] Paste	ANTHELCIDE [®] EQ Paste	QUEST [®] Gel <i>(If at least 6 months of age)</i> OR ANTHELCIDE EQ*	QUEST [®] PLUS Gel	QUEST Gel or STRONGID Paste**	
LATE FOALS BORN AFTER MARCH 	TARGET PARASITES	ROUNDWORMS (ascarids)	ROUNDWORMS (ascarids)	STRONGYLES & TAPEWORMS ROUNDWORMS* <i>(Only if detected via FEC)</i>	STRONGYLES	STRONGYLES	
	DEWORMER	STRONGID [®] Paste	ANTHELCIDE [®] EQ Paste	QUEST [®] PLUS Gel <i>(If at least 6 months of age)</i> OR ANTHELCIDE [®] EQ*	QUEST [®] Gel or STRONGID [®] Paste**	QUEST [®] Gel or ANTHELCIDE [®] EQ Paste**	

*At weaning, ask your veterinarian to conduct a fecal egg count (FEC) test to determine if a final roundworm treatment is needed between 6-9 months of age.

**If verified still effective against small strongyles through fecal egg count reduction test (FECRT).

Do not use QUEST Gel or QUEST PLUS Gel in foals less than 6 months of age or in sick, debilitated and underweight horses. Do not use in other animal species, as severe adverse reactions, including fatalities in dogs, may result. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

¹American Association of Equine Practitioners. Parasite Control Guidelines. <https://aaep.org/guidelines/parasite-control-guidelines>. Accessed March 1, 2021.

All trademarks are the property of Zoetis Services LLC or a related company or a licensor unless otherwise noted.

© 2021 Zoetis Services LLC. All rights reserved. QST-00171

