


Administering SILEO[®]

(dexmedetomidine oromucosal gel)




Wear Gloves

Wear impermeable disposable gloves when handling SILEO and handling the oral syringe.




Hold Plunger

Hold the oral syringe so that you can see the dot markings on the oral syringe plunger.




Unlock

Hold the plunger with your left hand and unlock the green ring-stop by turning it toward you until it is able to slide freely.




Move Ring

Move the ring-stop to the opposite end of the plunger.




Lock

Hold the plunger with your right hand and lock the ring-stop by turning it away from you.




Unlock

Hold the plunger with your right hand and unlock the ring-stop by turning it toward you. **Do not pull the plunger!**




Move Ring

Move the ring-stop toward the other end of the plunger for choosing the correct dose based on your veterinarian's prescription.



Set Dose and Lock

Position the ring-stop so that the side nearest the barrel is in line with the graduation mark (black line) and the required number of dots shows between the ring-stop and the barrel. Lock the ring-stop by turning it away from you. **Before dosing, make sure that the ring-stop is locked.**




Pull Cap (Tight)

Pull the cap strongly while holding the barrel. **Note:** The cap is very tight (pull, do not twist). Save the cap for replacement.




Dose Into Cheek

Place the oral syringe tip between the dog's cheek and gum and press the plunger until the ring-stop causes the plunger to stop.



IMPORTANT

The gel should not be swallowed. If the gel is swallowed, it may not be effective. Feeding and giving treats within 15 minutes after administration should be avoided.



Back to Package

Re-cap the oral syringe and return it to the outer package as the product is sensitive to light. Make sure that the carton is closed properly.

Keep the package out of sight and reach of children at all times. Remove and discard gloves. Use or discard syringe contents within 4 weeks after opening the syringe. To discard the product, please return your carton with the syringe back to your veterinarian.