
# PRODUCT PROFILE MG-BAC®



# **DESCRIPTION**

MG-Bac® is an inactivated vaccine for the immunization of healthy chickens and turkeys as an aid in the prevention of clinical signs associated with *Mycoplasma gallisepticum* infection.

#### COMPOSITION

Inactivated M. gallisepticum

# **FORMULATION**

Water-in-oil emulsion Option: Ready to use

#### **TARGET SPECIES**

Chickens and turkeys

#### **INDICATIONS**

MG-Bac is recommended for vaccination of healthy chickens and healthy turkeys 1 week of age or older to help protect against clinical signs associated with *M. gallisepticum* infection.



#### **VACCINATION PROGRAM**

- Inject healthy chickens or turkeys 1 to 10 weeks of age subcutaneously (in the lower neck region) using aseptic technique. For birds 10 weeks of age or older, one may vaccinate intramuscularly or subcutaneously.
- For optimum results, vaccinate twice, allowing at least
  4 weeks between vaccinations. The vaccination program
  should be completed prior to the point of lay.
- Revaccinate birds during molt before the second laying period.
- Dosage: One dose (0.5 mL) per bird.

# **METHOD OF ADMINISTRATION**

Subcutaneous injection (in the lower neck region), or intramuscularly (only in birds 10 weeks of age or older).

## WITHHOLDING PERIOD

42 days before slaughter

# **PRESENTATION**

1 x 500 mL (1,000 doses) bottles

## **STORAGE**

- Store in the dark at 36°F to 45°F (2°C to 7°C).
- Do not freeze.



# **PRECAUTIONS**

- Vaccinate only healthy birds prior to field exposure.
- If birds are vaccinated during lay, a drop in egg production may occur.
- Warm vaccine to 72°F (22°C) and shake well before using.
- Use entire contents when first opened.
- Penicillin added as a preservative.

# **CONTRAINDICATIONS**

Do not vaccinate broilers or meat turkeys by the intramuscular route of administration.

# **WARNINGS**

In case of accidental human injection, seek immediate medical attention, stating the vaccine is an oil-emulsion type.

# GET RELIABLE PROTECTION AND REWARDING RESULTS. GET MG-BAC®

Contact your local Zoetis representative for more information.

Size*	Standard Pack
500 mL 1,000 doses	1



**Zoetis U.S. Poultry Headquarters** 1040 Swabia Court

Durham, NC 27703-8481 Phone: 800-849-3372

Customer Service: 855-702-1036

zoetisUS.com/poultry

