

Strongid[®] C

(pyrantel tartrate)

Equine Anthelmintic

Medicated For Continuous Feeding

Net Weight: 25 lb (11.3 kg)

For the prevention of *Strongylus vulgaris* larval infestation in horses

For control of the following parasites in horses:

Large Strongyles (adults): *S. vulgaris*, *S. edentatus*

Small Strongyles (adults and 4th-stage larvae): *Cyathostomum* spp., *Cylicocyclus* spp., *Cylicostephanus* spp., *Cylicodontophorus* spp., *Poteriostomum* spp., *Triodontophorus* spp.

Pinworms (adults and 4th-stage larvae): *Oxyuris equi*

Ascarids (adults and 4th-stage larvae): *Parascaris equorum*

Active Drug Ingredient:

pyrantel tartrate..... 1.06% (4.8 g/lb)

Guaranteed Analysis:

Crude Protein, not less than 12.5%

Crude Fat, not less than 2.0%

Crude Fiber, not more than 22.0%

Ingredients:

Dehydrated Alfalfa Meal, Wheat Middlings, Ground Corn, Cane Molasses, Mineral Oil, preserved with propionic acid.

Mixing and Feeding Directions

Strongid C is to be administered on a continuous basis either as a top-dress or mixed in the horse's daily grain ration at the rate of 1.2 mg pyrantel tartrate per lb of body weight daily. To achieve this dose, administer 1 oz of Strongid C per 250 lb of body weight. (A Strongid C measuring cup is enclosed.) Strongid C should be administered

for the entire period that the animal is at risk to internal parasites. Unprotected animals that have grazed may already have an established *S. vulgaris* larval infestation. Before administering Strongid C, these animals should be treated with a therapeutic dose of a larvicidal product.

Foals may be administered Strongid C as soon as consistent intake of grain mix is occurring. This is generally between 2-3 months of age.

Strongid C may be used in mares at any stage of pregnancy or lactation. Stallion fertility is not affected by the use of Strongid C.

Top-Dress Directions	
Body Weight (lb)	oz per Day of Strongid C
250	1
500	2
750	3
1000	4
1250	5

(Measuring Cup Enclosed)

Medicated Grain Mix Directions			
lb of Medicated Grain mix per 100 lb of Body Weight	lb of Strongid C	lb of Non-medicated Feed	Concentration Grams/Ton
2.0	25	1975	120
1.5	33	1967	160
1.0	50	1950	240
0.5	100	1900	480
0.2	250	1750	1200


Caution: Consult your veterinarian before using in severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism. Do not mix in feeds containing bentonite.

⚠ **Warning:** Not for use in horses intended for human consumption. ⬅



zoetis

Distributed by:
Zoetis Inc.
Kalamazoo, MI 49007



10006000
Made in USA