

Long-lasting relief from canine allergic and atopic dermatitis is now a breeze!

CYTOPOINT®



How to dose

- Cytopoint® is available in single-use 1 mL vials in 4 concentrations: **10 mg, 20 mg, 30 mg or 40 mg/vial**
- Administer Cytopoint at a minimum dose of 2 mg/kg body weight. For convenience, the dosing table below may be used as a guideline
- Repeat administration every 4 to 8 weeks as needed in individual patients

DOG BODY WEIGHT		VIALS NEEDED				
	Pounds	10 mg	20 mg	30 mg	40 mg	TOTAL VIALS
DOGS <5 lb (<2.3 kg) 0.09 mL/lb (0.2 mL/kg) drawn from ONE 10-mg vial	<5	<1				<1
DOGS 5-40 lb (2.3-18.1 kg) Draw full volume from ONE 1-mL vial as indicated	5.0 - 10	1				1
	10.1 - 20		1			1
	20.1 - 30			1		1
	30.1 - 40				1	1
DOGS >40 lb (>18.1 kg) Requires full volume of TWO OR MORE 1 mL vials as indicated Draw the entire dose (or volume) from each vial into 1 syringe and administer as a single injection	40.1 - 50	1			1	2
	50.1 - 60		1		1	2
	60.1 - 70			1	1	2
	70.1 - 80				2	2
	80.1 - 90	1			2	3
	90.1 - 100		1		2	3
	100.1 - 110			1	2	3
	110.1 - 120				3	3
	120.1 - 130	1			3	4
	130.1 - 140		1		3	4
	140.1 - 150			1	3	4
	150.1 - 160				4	4
	160.1 - 170	1			4	5
	170.1 - 180		1		4	5
	180.1 - 190			1	4	5
	190.1 - 200				5	5



Store vials upright, in original packaging, at 2-8 °C/ 35.6-46.4 °F. Do not freeze.

– Prolonged exposure to a higher temperature and/or direct sunlight may adversely affect potency



Cytopoint contains no preservatives, and vials are single-use only



Vials should be discarded after puncture—do not split vials between patients

Indications: Cytopoint has been shown to be effective for the treatment of dogs against allergic dermatitis and atopic dermatitis.

All trademarks are the property of Zoetis Services LLC or a related company or a licensor unless otherwise noted.

© 2025 Zoetis Services LLC. All rights reserved. CYT-00235R3

zoetis