









BABIES NEED SPECIAL DEWORMING CARE

A TARGETED APPROACH FOR CONTROLLING PARASITES DEPENDS ON WHEN YOUR FOAL WAS BORN

EARLY FOALS BORN MARCH OR EARLIER	AGE	2 MONTHS	4-6 MONTHS (Prior to weaning)	7-9 MONTHS (Approx. 1 month post-weaning)	9 MONTHS	12 MONTHS
	TARGET PARASITES	ROUNDWORMS (ascarids)	ROUNDWORMS (ascarids)	STRONGYLES	STRONGYLES & TAPEWORMS	STRONGYLES
				ROUNDWORMS* (Only if detected via FEC)		
	DEWORMER	ANTHELCIDE® EQ Paste	STRONGID® Paste	QUEST® Gel (If at least 6 months of age)	QUEST® PLUS Gel	QUEST Gel or STRONGID Paste**
				ANTHELCIDE EQ*		

LATE FOALS BORN AFTER MARCH	AGE	2 MONTHS	4-6 MONTHS (Prior to weaning)	7-9 MONTHS (Approx. 1 month post-weaning)	9 MONTHS	12 MONTHS
Movay ovavovi	TARGET PARASITES	ROUNDWORMS (ascarids)	ROUNDWORMS (ascarids)	STRONGYLES & TAPEWORMS ROUNDWORMS* (Only if detected via FEC)	STRONGYLES	STRONGYLES
	DEWORMER	ANTHELCIDE® EQ Paste	STRONGID® Paste	QUEST® PLUS Gel (If at least 6 months of age) OR ANTHELCIDE® EQ*	QUEST® Gel or STRONGID® Paste**	QUEST® Gel or ANTHELCIDE® EQ Paste**

^{*}At weaning, ask your veterinarian to conduct a fecal egg count (FEC) test to determine if a final roundworm treatment is needed between 6-9 months of age.

Do not use QUEST Gel or QUEST PLUS Gel in foals less than 6 months of age or in sick, debilitated and underweight horses. Do not use in other animal species, as severe adverse reactions, including fatalities in dogs, may result. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.



^{**}If verified still effective against small strongyles through fecal egg count reduction test (FECRT).