


Dosing and Administration

Librela[™]
(bedinvetmab injection)



Concentrations:

5 mg, 10 mg, 15 mg, 20 mg, and 30 mg of bedinvetmab

Dose:

0.23 mg/lb (0.5 mg/kg) body weight

Dosing frequency:

Once monthly

Administration:

Subcutaneous injection

Storage:

Librela should be stored in a refrigerator between 36 °F to 46 °F (2 °C to 8 °C). Do not freeze. Store vials in their boxes to protect from prolonged exposure to light.

Once punctured, contents of the vial should be used immediately and any remaining solution should be discarded.

For use in dogs only.

Weight	Librela Strength				
	5 mg/mL	10 mg/mL	15 mg/mL	20 mg/mL	30 mg/mL
11-22.1 lb (5-10 kg)	1 Vial				
22.2-44.1 lb (10.1-20 kg)		1 Vial			
44.2-66.1 lb (20.1-30 kg)			1 Vial		
66.2-88.2 lb (30.1-40 kg)				1 Vial	
88.3-132.3 lb (40.1-60 kg)					1 Vial
132.4-176.4 lb (60.1-80 kg)				2 Vials	
176.5-220.5 lb (80.1-100 kg)				1 Vial	+ 1 Vial
220.6-264.6 lb (100.1-120 kg)					2 Vials

- **For dogs <11 lb (<5 kg):** Aseptically withdraw 0.045 mL/lb (0.1 mL/kg) from a 5 mg/mL vial into a single syringe and administer immediately. Discard the vial after the dose has been withdrawn

- **For dogs ≥11 lb (≥5 kg):** Dogs should be dosed by weight range according to the Dosing Chart above. Dogs are given the full content of 1 or 2 vials of the appropriate concentration based on body weight. Aseptically withdraw the total dose into a single syringe and administer immediately

For more information about Librela, visit **LibrelaVetTeam.com**

IMPORTANT SAFETY INFORMATION: For use in dogs only. Women who are pregnant, trying to conceive or breastfeeding should take extreme care to avoid self-injection. Hypersensitivity reactions, including anaphylaxis, could potentially occur with self-injection. LIBRELA should not be used in breeding dogs or in pregnant or lactating dogs. LIBRELA should not be administered to dogs with known hypersensitivity to bedinvetmab. The most common adverse events reported in a clinical study were urinary tract infections, bacterial skin infections and dermatitis. See full Prescribing Information at [LibrelaPI.com](https://www.librelapi.com).