


# Dosing and Administration

**Librela**<sup>™</sup>  
(bedinvetmab injection)



**Concentrations:**

5 mg, 10 mg, 15 mg, 20 mg, and 30 mg of bedinvetmab

**Dose:**

0.23 mg/lb (0.5 mg/kg) body weight

**Dosing frequency:**

Once monthly

**Administration:**

Subcutaneous injection

**Storage:**

Librela should be stored in a refrigerator between 36 °F to 46 °F (2 °C to 8 °C). Do not freeze. Store vials in their boxes to protect from prolonged exposure to light.

Once punctured, contents of the vial should be used immediately and any remaining solution should be discarded.

**For use in dogs only.**

Weight	Librela Strength				
	5 mg/mL	10 mg/mL	15 mg/mL	20 mg/mL	30 mg/mL
<b>11-22.1 lb</b> (5-10 kg)	<b>1 Vial</b>				
<b>22.2-44.1 lb</b> (10.1-20 kg)		<b>1 Vial</b>			
<b>44.2-66.1 lb</b> (20.1-30 kg)			<b>1 Vial</b>		
<b>66.2-88.2 lb</b> (30.1-40 kg)				<b>1 Vial</b>	
<b>88.3-132.3 lb</b> (40.1-60 kg)					<b>1 Vial</b>
<b>132.4-176.4 lb</b> (60.1-80 kg)				<b>2 Vials</b>	
<b>176.5-220.5 lb</b> (80.1-100 kg)				<b>1 Vial</b>	<b>+</b> <b>1 Vial</b>
<b>220.6-264.6 lb</b> (100.1-120 kg)					<b>2 Vials</b>

• **For dogs <11 lb (<5 kg):** Aseptically withdraw 0.045 mL/lb (0.1 mL/kg) from a 5 mg/mL vial into a single syringe and administer immediately. Discard the vial after the dose has been withdrawn

• **For dogs ≥11 lb (≥5 kg):** Dogs should be dosed by weight range according to the Dosing Chart above. Dogs are given the full content of 1 or 2 vials of the appropriate concentration based on body weight. Aseptically withdraw the total dose into a single syringe and administer immediately

For more information about Librela, visit [LibrelaVetTeam.com](http://LibrelaVetTeam.com)

**IMPORTANT SAFETY INFORMATION:** For use in dogs only. Women who are pregnant, trying to conceive or breastfeeding should take extreme care to avoid self-injection. Hypersensitivity reactions, including anaphylaxis, could potentially occur with self-injection. LIBRELA should not be used in breeding dogs or in pregnant or lactating dogs. LIBRELA should not be administered to dogs with known hypersensitivity to bedinvetmab. The most common adverse events reported in a clinical study were urinary tract infections, bacterial skin infections and dermatitis. See full Prescribing Information at [LibrelaPI.com](http://LibrelaPI.com).