Ask your veterinarian for Stablelab



DO YOU KNOW YOUR HORSE'S SAA?

Measurement of Serum Amyloid A (SAA) with Stablelab can detect infection before other clinical signs appear. In healthy horses, SAA levels in the blood are very low. With infectious conditions, SAA rapidly rises and stays elevated until the infection subsides. Stablelab can quickly and effectively assist your veterinarian in detecting infectious conditions and monitoring your horse's treatment response.^{13,4,5}



HIGH STANDARD OF CARE

Stablelab allows your veterinarian to perform a blood test in only 10 minutes to aid in the examination and care of your horse.



GENERAL WELLNESS BASELINE

Having an SAA value aids in the clinical detection of inflammation due to possible infection at the time of exam.



BIOSECURITY FOR INFECTIOUS DISEASE

Monitoring possible spread of disease where horses are housed can aid in the control and quaratine protocols. 7



MONITORING RESPONSE TO MEDICATION

Repeat testing allows for monitoring progression of effectiveness of therapy.

二 ふく
-~~_
0-0-0-

CRITICAL OR EMERGENCY CASES

When every second counts, measuring SAA at your horse's side, provides immediate, reliable results, and can expedite appropriate diagnosis and treatment.



DETECTING INFECTION IN FOALS

Newborn foals can decline rapidly so monitoring SAA values can identify a potential problem before it gets out of hand.⁶



POOR PERFORMANCE

For horses not performing up to expectations, measurement of SAA quickly helps to evaluate general health and rule out subclinical infectious conditions.²



PRE-AND POST-PERFORMANCE EVALUATION

Measurement of SAA can give you peace of mind that your horse is not brewing a subclinical infection that may impact performance prior to competition. Additionally, Stablelab can help screen for a possible infectious disease before returning home, that may affect stablemates.²



PRE-PURCHASE EXAMINATION

In addition to other routine screening bloodwork, incorporating SAA measurement using Stablelab can help identify subclinical infectious disease prior to purchase.



Enhance Your Horse's Clinical Exam





DIFFERENTIATE INFECTION FROM NON-INFECTION

With infectious inflammation, SAA almost always elevates above 50 μ g/mL, while non-infectious inflammatory conditions rarely elevate SAA at all.^{1,2,5,8} When the horse contracts an infection, SAA rapidly and dramatically rises and once the infection has resolved, SAA will drop by 50% every 24 hours allowing your veterinarian to manage a case with greater clarity than ever before.⁹



30 TIMES MORE SENSITIVE THAN A THERMOMETER

SAA was found to be more sensitive than a thermometer and better than traditional lab tests at detecting subclinical infections^{1,2}



AN EVOLUTION IN RAPID DIAGNOSTICS

Stablelab is a significant advancement in stall-side, point-of-care equine diagnostics. Stablelab is trusted by thousands of equine veterinarians in 25 countries worldwide.¹⁰

Measuring and monitoring Serum Amyloid A, in conjunction with other diagnostic tests, enables your veterinarian to **detect**, **monitor** and **screen** your horse.



Learn more at ZoetisEquine.com/Stablelab

REFERENCES

- ¹ Belgrave, R. et al. Assessment of Serum Amyloid A testing of horses and its clinical application in a specialized equine practice. JAVMA, Vol 243 no 1, July 1, 2013. Pages 113–119. ² Oertly et al. Early Inflammation in Horses Following Long-Distance Transportation by Air. AAEP Proceedings, Vol 62. Pages 460–461. 2017.
- ³ Viner, M et al. Comparison of Serum Amyloid A in horses with Infectious and noninfectious respiratory diseases. Journal of Equine Veterinary Science. 2017 (49) 11–13.
- ⁴ Jacobsen, Stine, Review of equine Acute-Phase proteins, AAEP Proceedings Vol 53 2007, 230-235.
- ⁵ Nolen–Walston, R. How to interpret Serum Amyloid A concentrations. AAEP Proceedings 2015, 130–137.
- ⁶ McCracken JL. Evaulation of the Use of White Blood Cell Count, Fibrinogren, Serum Amyloid A and Ultrasonographic Grade to Refine A Thoracic Rhodococcus equi Screening Program. In Proceedings. American Association of Equine Practitioners. 2019. 65; 522–530.
- ⁷ Pepys MB, Baltz ML, Tennent GA. Serum Amyloid A (SAA) in horses: Objective measurement of the Acute Phase Response. Equine Veterinary Journal. 1989. 21(2); 106–109.
- ⁶ Anhold et. al. A comparison of Elevated Blood Parameter Values in a Population of Thoroughbred Racehorses. JEVS 34 (2014) 651–655
- [°] Ludwig EK, Wiese RB, Graham MR et al. Serum and Synovial Fluid Serum Amyloid A Response in Equine Models for Synovitis and Septic Arthritis. Veterinary Surgery 46 (2016) 859–867.
- ¹⁰ Customer sales data on file.

All trademarks are the property of Zoetis Services LLC or a related company or a licensor unless otherwise noted. Stablelab is a registered trademark of Epona Biotech Limited, used under license. © 2023 Zoetis Services LLC. All rights reserved. STB-00025R2

