

# CONVENIA® (cefovecin sodium) in your practice



## CONVENIA Dosing Chart at 3.6 mg/lb (8 mg/kg or 0.045 mL/lb)

Dog Weight	CONVENIA Volume
5 lb	0.23 mL
10 lb	0.45 mL
15 lb	0.67 mL
20 lb	0.90 mL
30 lb	1.40 mL
40 lb	1.80 mL
50 lb	2.30 mL
60 lb	2.70 mL
70 lb	3.20 mL
80 lb	3.60 mL
90 lb	4.10 mL
100 lb	4.50 mL



- Label dose for dogs and cats = 3.6 mg/lb (8 mg/kg or 0.045 mL/lb) subcutaneously
- Reconstituted concentration = 80 mg/mL
- Reconstitute with 10 mL of sterile water for injection, USP
- Use within 56 days after reconstitution
- Always refrigerate in original carton to protect from light
- Minimum pet age for use is 4 months



CONVENIA will naturally turn amber over time after reconstitution. The color change has no impact on safety or efficacy, provided CONVENIA is kept under proper storage conditions.

**IMPORTANT SAFETY INFORMATION:** People with known hypersensitivity to penicillin or cephalosporins should avoid exposure to CONVENIA. Do not use in dogs or cats with a history of allergic reactions to penicillins or cephalosporins. Side effects for both dogs and cats include vomiting, diarrhea, decreased appetite/anorexia and lethargy. Please see attached full Prescribing Information.

