

USING MEDICATED FEED ADDITIVES FROM ZOETIS WITH THE VETERINARY FEED DIRECTIVE

AUREOMIX® S, AUREOMYCIN® and LINCOMIX® each help swine veterinarians and producers reliably treat bacterial diseases. Veterinary Feed Directive (VFD) regulations require that these feed additives involve a veterinarian to oversee correct use of these products in operations.

Producers need to have a valid veterinary client patient relationship (VCPR) as defined by state regulation in order for veterinarians to issue a VFD.

Aureomix® S

Granular

WHAT DOES VFD MEAN FOR HOW I USE AUREOMIX S 10/10 AND AUREOMIX S 40/40?

PRODUCERS: Producers must request and obtain a VFD from their veterinarian to feed rations with AUREOMIX S to swine.

FEED DISTRIBUTORS: Those who handle products or mix feed with AUREOMIX S need to file a one-time notice (Letter of Distribution) with the Food and Drug Administration (FDA) declaring they intend to distribute feed or product with VFD medicated feed additives.

CLAIM:

- **100 g/ton of AUREOMYCIN brand chlortetracycline and 100 g/ton sulfamethazine:** For reduction of the incidence of cervical abscesses; treatment of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis* and vibronic dysentery); prevention of these diseases during times of stress; and maintenance of weight gains in the presence of atrophic rhinitis. Feed as the sole ration. Withdraw 15 days prior to slaughter.

Aureomycin®

WHAT DOES VFD MEAN FOR HOW I USE AUREOMYCIN?

PRODUCERS: Producers must request and obtain a VFD from their veterinarian to feed rations with AUREOMYCIN to swine.

FEED DISTRIBUTORS: Those who handle products or mix feed with AUREOMYCIN need to file a one-time notice (Letter of Distribution) with the FDA declaring they intend to distribute feed or product with VFD medicated feed additives.

CLAIMS:

- **10 mg per lb. of body weight per day:** Control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis* susceptible to chlortetracycline. Feed continuously for not more than 14 days.
- **10 mg per lb. of body weight per day:** Treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to chlortetracycline. Feed continuously for not more than 14 days.
- **50 to 100 g/ton:** For reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by Group E *streptococci* susceptible to chlortetracycline.
- **400 g/ton:** Breeding swine: For the control of leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by *Leptospira pomona* susceptible to chlortetracycline. Feed continuously for not more than 14 days.



Lincomix®

WHAT DOES VFD MEAN FOR HOW I USE LINCOMIX?

PRODUCERS: Producers must request and obtain a VFD from their veterinarian to feed rations with LINCOMIX® to swine.

FEED DISTRIBUTORS: Those who handle products or mix feed with LINCOMIX need to file a one-time notice (Letter of Distribution) with the FDA declaring they intend to distribute feed or product with VFD medicated feed additives.

CLAIMS:

- **40 g/ton:** For control of swine dysentery and the control of porcine proliferative enteropathy (ileitis) caused by *Lawsonia intracellularis*. For use in animals or on premises with a history of swine dysentery, but where symptoms have not yet occurred. Feed as the sole ration.
- **100 g/ton:** For treatment of swine dysentery and the control of porcine proliferative enteropathy (ileitis) caused by *Lawsonia intracellularis*. Feed as sole ration for 3 weeks or until signs of disease (watery, mucoid or bloody stools) disappear.
- **100 g/ton followed by 40 g/ton:** For the treatment and control of swine dysentery and the control of porcine proliferative enteropathy (ileitis) caused by *Lawsonia intracellularis*. Feed 100 g/ton of complete feed as the sole ration for 3 weeks or until signs of disease (watery, mucoid or bloody stools) disappear, followed by 40 g/ton of lincomycin.
- **100-200 g/ton:** For the reduction in the severity of respiratory disease associated with *Mycoplasma hyopneumoniae*. Feed as sole ration for 21 days.

QUICK FACTS ABOUT VFD:

- ✓ A Veterinary Feed Directive (VFD) is required for producers to include AUREOMIX® S, AUREOMYCIN® and LINCOMIX® in rations.
- ✓ Producers need to have a valid VCPR in order for veterinarians to issue a VFD.
- ✓ A VFD is specific to client, group of animals, indications and time. For more about specific indications for AUREOMIX S, AUREOMYCIN and LINCOMIX, visit zoetisUS.com.
- ✓ A VFD can be issued for approved indications only. No extra-label use of medicated feed additives is allowed. For more about specific indications for AUREOMIX S, AUREOMYCIN and LINCOMIX, visit zoetisUS.com.
- ✓ A VFD can be issued for a maximum of six months.
- ✓ The producer, feed distributor and veterinarian must keep a copy of the VFD on file for two years.
- ✓ The veterinarian must be licensed to practice in the state where the pigs will be fed.

WE ARE COMMITTED TO SUPPORTING YOU IN THE RESPONSIBLE USE OF ANTIBIOTICS. FOR MORE DETAILS ABOUT THE CHANGES IMPLEMENTED ON JAN. 1, 2017, CONTACT YOUR ZOETIS REPRESENTATIVE OR VISIT RESPONSIBLEANTIBIOTICUSE.COM.

Withdraw AUREOMIX S fifteen (15) days prior to slaughter.

Do not use LINCOMIX in swine intended for breeding. Do not allow unapproved species access to feeds containing lincomycin.

All trademarks are the property of Zoetis Services LLC or a related company or a licensor unless otherwise noted.
© 2017 Zoetis Services LLC. All rights reserved. AMS-00011

zoetis